



FAU Institutional Repository

<http://purl.fcla.edu/fau/fauir>

This paper was submitted by the faculty of [FAU's Harbor Branch Oceanographic Institute](#).

Notice: ©2005 American Chemical Society and American Society of Pharmacognosy. This document is the accepted manuscript version of a published work that appeared in final form in *Journal of Natural Products* after peer review and technical editing by the publisher. To access the final edited and published work see <http://dx.doi.org/10.1021/np040147u>. This article may be cited as: Reddy, N. S., Reed, J. K., Longley, R. E., & Wright, A. E. (2005). Two New Cytotoxic Linderazulenes from a Deep-Sea Gorgonian of the Genus *Paramuricea*. *Journal of Natural Products*, 68(2), 248-250.

Two New Cytotoxic Linderazulenes from a Deep-Sea Gorgonian of the Genus *Paramuricea*


N. Srinivasa Reddy, John K. Reed, Ross E. Longley,[†] and Amy E. Wright*

Division of Biomedical Marine Research, Harbor Branch Oceanographic Institution, 5600 US 1 North, Fort Pierce, Florida 34946

Received June 30, 2004

The known compound linderazulene (**1**) and two new linderazulenes (**2**, **3**) were isolated from a deep-sea gorgonian *Paramuricea* sp. The structures of **2** and **3** were determined through spectroscopic methods. Compounds **1–3** show moderate in vitro cytotoxicity against the P388 murine leukemia cell line with IC₅₀'s of 18.8, 2.7, and 15.6 μg/mL, respectively. Compound **2** showed moderate activity against the PANC-1 pancreatic cell line with an IC₅₀ of 18.7 μg/mL.

Gorgonians have proven to be a rich source of guaiazulene-related pigments.¹ These compounds show a variety of activities including antifungal, antitumor, antibacterial, immunoregulatory, and antiproliferative effects on fertilized sea urchin and ascidian eggs.^{2–7} As part of our continued research on biologically active secondary metabolites from marine organisms,⁸ a deep-sea gorgonian of the genus *Paramuricea* (family Paramuriceidae) collected at Westpunt, Curaçao, was investigated. The EtOAc partition of the gorgonian was fractionated by regular-phase vacuum column chromatography followed by reverse-phase HPLC to yield three compounds (**1**, **2**, and **3**). Compound **1** was identified as the known compound linderazulene by comparison of its spectroscopic data to the published data. Linderazulene was first reported from the gorgonians *Acalycigorgia* sp.⁴ and *Paramuricea chamaeleon*.⁹ Detailed interpretation of the spectroscopic data observed for **2** and **3** identified them as new members of the linderazulene class of compounds. This paper describes the isolation, structure elucidation, and cytotoxic activity of compounds **2** and **3**.



Compound **2** was obtained as a pink amorphous solid (mp 138–139 °C). HRFABMS suggested a formula of C₁₆H₁₄O₃ for **2** (*m/z* 255.1033 [M + H]⁺ observed, 255.1021 calculated) requiring 10 degrees of unsaturation. This formula was further supported by the ¹³C, DEPT, and HMQC spectra (see Table 1), which indicated the presence of 16 carbon resonances including seven quaternary olefinic

carbons (δ 159.0, 141.1, 139.3, 136.3, 126.3, 120.3, 116.4), five olefinic methine carbons (δ 141.1, 137.0, 130.7, 116.1, 115.7) two methyl carbons (δ 25.2, 8.0), one methoxyl carbon (δ 50.9), and one ester carbon (δ 166.3). The ¹H NMR spectrum showed resonances corresponding to two aromatic methyls (δ 2.45 (s, 3H) and 2.93 (s, 3H)); one ester methyl (δ 3.94 (s, 3H)); two mutually coupled olefinic resonances (δ 8.24 (d, *J* = 3.8 Hz, 1H) and 7.28 (d, *J* = 3.8 Hz, 1H)); and three one-proton aromatic singlets (δ 9.99, 7.69, and 7.56). The IR spectrum of compound **2** showed bands corresponding to an ester carbonyl functionality (1678 cm⁻¹). The foregoing spectral data suggested that compound **2** is a new linderazulene metabolite in which the C-11 methyl has been oxidized to an ester. Analysis of the ¹H–¹H COSY and proton-detected HMQC and HMBC spectra allowed for full assignment of the NMR data (Table 1). An alternative structure in which the C-11 ester functionality is attached at C-7 could be ruled out on the basis of correlations observed in the NOESY spectrum between H-7 and H₃-12 as well as by chemical shift arguments. We propose the common name 11-carbomethoxy-linderazulene for **2**.

Compound **3** was also obtained as an amorphous pink solid (mp 136 °C). HRFABMS suggested a formula of C₁₅H₁₂O₂ for **3** (*m/z* 225.0917 [M + H]⁺ observed, 225.0915 calculated) requiring 10 degrees of unsaturation. The NMR spectra of **3** (Table 1) were nearly identical to those observed for **2** with the primary difference being the observation of resonances attributable to an aldehyde functionality [¹³C NMR: δ 187.1 (CH); ¹H NMR: δ 10.3 (s, 1H)] rather than the ester functionality observed for **2**. The IR spectrum of compound **3** confirmed the presence of an aromatic aldehyde functionality (ν 2746, 1641 cm⁻¹). A detailed analysis of the 2D spectra (COSY, NOESY, HMQC, and HMBC) confirmed the assignment of **3** as 11-formyl-linderazulene.

Compounds **1–3** show moderate in vitro cytotoxicity against the P388 murine leukemia cell line with IC₅₀'s of 18.8, 2.7, and 15.6 μg/mL, respectively. Compound **2** also showed moderate activity against the PANC-1 pancreatic tumor cell line with an IC₅₀ of 18.7 μg/mL. When tested for their ability to inhibit the growth of the fungal pathogen *Candida albicans*, no activity was detected at a concentration of 50 μg/mL under the conditions tested.⁸

Experimental Section

General Experimental Procedures. Melting points were recorded on a Gallenhampt melting point apparatus; IR spectra were collected on a MIDAC M-1200 with Galactac GRAMS/

* Corresponding author. Phone: (772) 465-2400, ext. 459. Fax: (772) 461-2221. E-mail: wright@hboi.edu.

[†] Current address: Taxolog Inc., 3216 Sessions Rd., Suite 100, Tallahassee, FL 32303.

Table 1. NMR Spectral Data for Compounds **2** and **3** in CDCl₃

atom no.	compound 2				compound 3		
	δ_C , DEPT	δ_H , mult. (<i>J</i> in Hz)	HMBC ^a	NOESY	δ_C , DEPT	δ_H , mult. (<i>J</i> in Hz)	HMBC ^a
2	141.1 CH	7.56 s	C-3, C-3a, C-9a	H-10	141.8 CH	7.56 s	C-3, C-3a, C-9a
3	120.3 C				120.4 C		
3a	126.3 C				126.6 C		
4	130.7 CH	9.99 s	C-3a, C-4a, C-7a, C-9a	H-10	131.5 CH	9.99 s	C-3a, C-4a, C-7a, C-9a
4a	116.4 C				117.4 C		
5	136.3 C				135.1 C		
6	137.0 CH	8.24 d (3.8)	C-4a, C-5, C-7a	H-7	141.3 CH	8.09 d (3.8)	C-4a, C-5, C-7a
7	116.1 CH	7.28 d (3.8)	C-4a, C-5, C-6, C-7a	H-6 H-12	117.5 CH	7.32 d (3.8)	C-4a, C-5, C-6, C-7a
7a	139.3 C				141.7 C		
8	141.1 C				141.5 C		
9	115.7 CH	7.69 s	C-3a, C-7a, C-8, C-9a, C-12	H-12	117.4 CH	7.80 s	C-3a, C-7a, C-8, C-9a, C-12
9a	159.0 C				159.4 C		
10	8.0 CH ₃	2.45 s 3H	C-2, C-3, C-3a	H-4, H-2	8.0 CH ₃	2.46 s 3H	C-2, C-3, C-3a
11	166.3 C				187.1 CH	10.29 s	
12	25.2 CH ₃	2.93 s 3H	C-7a, C-8, C-9	H-7, H-9	25.2 CH ₃	2.94 s 3H	C-7a, C-8, C-9
-OCH ₃	50.9 CH ₃	3.94 s 3H	C-11				

^a HMBC correlation from H number to carbon atoms listed.

386 software. The ¹H, COSY, NOESY, ¹³C, DEPT 135, DEPT 90, HMQC, and HMBC (optimized for 10 Hz) spectra were recorded on a Bruker AMX-500 operating at 500 MHz (¹H) and 125 MHz (¹³C). ¹H chemical shifts are referenced to CDCl₃ observed at 7.24 ppm, while ¹³C chemical shifts are referenced to CDCl₃ observed at 77.0 ppm. The HRFABMS and EIMS were obtained on a Kratos MS50TC mass spectrometer at Oregon State University.

Animal Material. The gorgonian specimen (HBOI ID 19-V-00-3-003) was collected May 19, 2000, using Harbor Branch Oceanographic Institution's *Johnson-Sea-Link* research submersible off the northwest coast of Curaçao, 12°22.78' N latitude and 69°10.15' W longitude, on a rocky slope and at a depth of 342 m. The specimen was frozen immediately after collection and kept frozen until workup. This specimen most closely fits the genus *Paramuricea* Kölliker, 1865 [Cnidaria, Anthozoa, Gorgonacea, Plexauridae (Paramuriceidae)], but cannot be ascribed to a known species at this time. It differs from other specimens that were collected in the same region, in both morphology and spiculation. This holaxonian gorgonian specimen was 40 cm tall, 30 cm wide, and branching mostly in one plane. The base of the main stem is 6 mm diameter, which has several secondary branches and numerous short (1–2 cm) terminal branchlets, 0.5–0.75 mm diameter, which do not anastomose. The polyps are distributed over the entire surface, tend to alternate about 1–2 mm apart, and are denser on the branchlets. The calyces (anthostele) are tubular to conical, ~0.5 mm high, and consist of relatively few thorn scale spicules with very blunt, slightly spinous, projecting spine and sparse basal root. The large anthocodia are exert (1–2 mm), with long tentacles and pinnules. The anthocodial collaret is relatively strong with a crown of 3 or 4 rows of thin, curved, slightly spinous spindles and some spinose rods. The coenenchyme consists of a thin layer of spicules, dominated by knee-shaped rods that are warty on the convex side and smooth on the concave side, and a few crosses. The axis is soft and fibrous with a cross-chambered central core. The coenenchymal spicules range from ~0.07 to 0.20 mm in length; the thorn scales are ~0.25 mm; the anthocodial spindles are ~0.15 to 0.45 mm; and the anthocodial rods are ~0.23 to 0.30 mm. The color of the specimen was described in situ from the submersible as pale lavender. While still fresh in the lab, the main stalk and branches were pale lavender, and the branchlets were tan. In alcohol the specimen is dark brown. A museum voucher specimen is deposited at the Harbor Branch Oceanographic Museum, catalog number 012:00804.

Extraction and Isolation. The frozen specimen of the gorgonian (100 g wet wt) was chopped into small slices and extracted exhaustively with EtOH (3 × 400 mL). The combined EtOH extracts were concentrated by distillation under reduced pressure. The resulting residue was partitioned between water

and EtOAc (3 × 100 mL). The combined EtOAc partition was concentrated to yield 1.2 g of a reddish brown solid. The EtOAc partition was fractionated by vacuum column chromatography on a Kieselgel 60H stationary phase using a step gradient of EtOAc in heptane. The fractions that eluted with heptane and heptane–EtOAc (95:5) (v/v) were combined and chromatographed by reversed-phase HPLC [Vydac C-18 Protein and Peptide column, 10 × 250 mm, H₂O–CH₃CN, 2:8, flow rate = 3 mL/min, detected by UV absorbance at 254 nm] to yield compounds **1** (1.4 mg, 0.0014%), **2** (21 mg, 0.021% of wet weight), and **3** (1.2 mg, 0.0012%).

11-Carbomethoxylinderazulene (2): amorphous pink solid, mp 138–139 °C; UV (MeOH) λ_{\max} (log ϵ) 389 (3.92), 376 (3.80), 325 (4.07), 308 (4.3), 247 (4.00), 228 (4.05) nm; IR (neat) ν_{\max} 2946, 2918, 2853, 1678, 1447, 1408, 1387, 1305, 1217, 1188, 1133, 1079, 1056, 939 cm⁻¹; ¹H and ¹³C NMR (CDCl₃), see Table 1; HRFABMS *m/z* 255.1033 (calcd for C₁₆H₁₅O₃ [M + H]⁺, 255.1021).

11-Formyllinderazulene (3): amorphous pink solid, mp 136 °C; UV (MeOH) λ_{\max} (ϵ) 395 (3.86), 332 (3.97), 316 (4.12), 293 (4.03), 260 (4.15), 215 (4.08) nm; IR (neat) ν_{\max} 2956, 2923, 2853, 2746, 1641, 1634, 1398, 1374, 1297, 1286, 1233, 1148, 1081, 1044, 943 cm⁻¹; ¹H and ¹³C NMR (CDCl₃), see Table 1; HRFABMS *m/z* 225.0917 (calcd for C₁₅H₁₃O₂ [M + H]⁺, 225.0915).

Biological Assays. P388 murine leukemia cells and PANC-1 pancreatic cancer cells were obtained from the American Type Culture Collection (Rockville, MD). Cytotoxicity assays were run as previously described according to the methods of Alley et al.¹⁰ Antifungal assays were run as previously described.⁸

Acknowledgment. One of the authors, N.S.R., is thankful to Harbor Branch Oceanographic Institution for a Postdoctoral Fellowship. Equipment funded by NIH grant number 2U19-CA 529955 was used during the course of the work described in this paper. We thank P. McCarthy and D. Harmody for antifungal testing. We thank J. Morre of Oregon State University, Corvallis, OR, for mass spectral data. This is Harbor Branch Oceanographic Institution Contribution number 1571 and The Center of Excellence in Biomedical and Marine Biotechnology Contribution number P200411. We thank Dr. A. Debrot, subdirector of the Caribbean Marine Biological Institute (CARMABI Foundation), Curacao, who provided exceptional assistance in planning and collaboration during this expedition. The success of this research cruise is due in large part to the CARMABI Foundation and the cooperation of Dr. Debrot and Dr. W. L. Bakhuis, Director of CARMABI. The scientific research and collections were authorized by the Netherlands Ministry of Foreign Affairs (Diplomatic Note VADV-172/00, January 31, 2000).

References and Notes

- (1) Faulkner, D. *J. Nat. Prod. Rep.* **2001**, *18*, 1–49, and previous reports in this series.
- (2) Fusetani, N.; Matsunaga, S.; Konosu, S. *Experientia* **1981**, *37*, 680–681.
- (3) Okuda, R. K.; Klein, D.; Kinnel, R. B.; Li, M.; Scheuer, P. J. *Pure Appl. Chem.* **1982**, *54*, 1907–1914.
- (4) Sakemi, S.; Higa, T. *Experientia* **1987**, *43*, 624–625.
- (5) Tanaka, J.; Miki, H.; Higa, T. *J. Nat. Prod.* **1992**, *55*, 1522–1524.
- (6) Ochi, M.; Kataoka, K.; Tatsukawa, A.; Kotsuki, H.; Shibata, K. *Chem. Lett.* **1993**, 2003–2006.
- (7) Seo, Y.; Rho, J. R.; Geum, N.; Yoon, J. B.; Shin, J. *J. Nat. Prod.* **1996**, *59*, 985–986.
- (8) Chen, Y.; Brian Killday, K.; McCarthy, P. J.; Schimoler, R.; Chilson, K.; Selitrennikoff, C.; Pomponi, S. A.; Wright, A. E. *J. Nat. Prod.* **2001**, *64*, 262–264.
- (9) Imre, S.; Thomson, R. H.; Yalhi, B. *Experientia* **1981**, *37*, 442–443.
- (10) Alley, M. C.; Scudiero, D. A.; Monks, A.; Hursey, M. L.; Czerwinski, M. J.; Fine, D. L.; Abbott, B. J.; Mayo, J. G.; Shoemaker, R. H.; Boyd, M. R. *Cancer Res.* **1988**, *48*, 589–596.

NP040147U